

Sampling For Nematodes

DR PAUL VINCELLI

LEXINGTON, KY.

April 24, 2012 – Nematodes are microscopic roundworms that attack roots of corn and other plants. In corn, nematodes affect yield by damaging the root system and interacting with opportunistic root-rotting fungi. They become more important when there are other root-related stresses in the same field, such as moisture shortage, root damage from insects, nutrient deficiencies, or compaction.

Although there is increasing interest in nematodes as potential yield-limiting factors, the threat these pose varies greatly from one field to the next. The only way to determine whether nematodes may be affecting yield is to sample for nematodes.

When to Sample

The window at 4-6 weeks after planting is probably the best time to sample, since the nematodes will be in the root zone and the soil is often moist enough to be able to insert a soil probe. Plus, sampling early in the growing season will give an idea as to whether nematodes are likely to affect crop productivity. Summer sampling is less than ideal. During summer, certain nematodes (like the sting nematode and needle nematode) move downward in the soil profile, where the moisture is located. Also, pulling samples from dry soil is difficult. Sampling after harvest has its advantages, since it gives a look at certain nematode populations, especially endoparasites (lance nematode and root-lesion nematode, which burrow within the corn root), as well as the nematodes that go deep into the soil during summer.

How to sample

The most important thing to realize is that sampling for corn nematodes is different than sampling for soybean cyst nematode. For corn, sampling both soil and roots is advisable for best results. Sampling guidelines are as follows:

Sampling soil

- Instead of sampling at random or zig-zag pattern (like you would for soybean cyst nematode), sample parts of the field where yield losses or symptoms (stunting, yellowing in elongated areas of the field) are not explained by other factors, such as soil compaction, soil type, etc. Corn nematode populations can be extremely variable (see Figure 1), so focus sampling in problem areas of the field. However, if some of the corn is severely affected, sample from the edge of the damaged areas rather than in the worst areas. It is important to avoid badly damaged corn, because corn nematodes require live roots to feed on, and if plants are severely damaged or dead, the nematode numbers will be low.

- Make sure to sample within the row – that's right, within the row. This is differ-

ent from sampling for soil fertility. Take 20-25 samples.

- Sampling for nematodes in corn requires a depth of at least 12 inches. This is much deeper than for soybean cyst nematode. If sampling in summer or fall, one may need to go as deep as 24-36 inches in order to detect sting nematode or needle nematode (although these two nematodes are usually found only in sandy soils).

- Put all 20-25 soil cores in a Ziploc bag. Don't break the soil cores, since some nematodes like stubby root nematode are extremely sensitive to soil disturbance. Label the bag with a permanent marker (not a magic marker, which comes off in water). Store samples in a cool, dry place until shipment.

Sampling roots

Collect five to ten root masses from V6 plants; the tops of the plants can be cut off and discarded, and soil adhering to roots can be removed as well. Roots can be placed in a plastic bag separate from your soil samples.

Where to send samples

Laboratories that can analyze nematodes in soil samples include:

- University of Illinois, <http://web.extension.illinois.edu/plantclinic/>
- Purdue University, <http://extension.entm.purdue.edu/nematology/cv/submissionform.pdf>
- Mississippi State University, <http://msucares.com/pubs/misc/m1230.pdf>
- University of Florida, <http://edis.ifas.ufl.edu/sr011>
- Iowa State University, <http://www.extension.iastate.edu/Publications/PD32.pdf>

DR. PAUL VINCELLI: Extension Professor and Provost's Distinguished Service Professor, University of Kentucky

2010 Corn Nematode Research Arenzville, IL

